

# Idaho, 2013

## Primary and secondary syphilis

- CASES: Idaho reported 15 cases in 2013.
- RATES: Idaho ranked 45 among 50 states reporting at least one case of P&S syphilis with 0.9 cases per 100,000 population compared to the U.S. rate of 5.5.

## Rates of primary and secondary syphilis, by sex

- In Idaho, the rate among males was 1.9 cases per 100,000 population compared to the U.S. male rate of 10.3.
- The rate among females was 0.0 compared to the U.S. female rate of 0.9.


## Rates of primary and secondary syphilis, by race/ethnicity

- In Idaho, the race/ethnicity adjusted rates per 100,000 population were 0.8 among whites, 0.0 among blacks, 2.2 among Hispanics, 0.0 among Asians, 0.0 among Native Hawaiian/Other Pacific Islanders, and 0.0 among American Indians/Alaska Natives.
- The rate among Hispanics was 2.6 times that of whites.


## Congenital syphilis

- CASES: Idaho reported 0 cases in 2013.
- RATES: Idaho had a rate of 0.0 cases per 100,000 live births compared to the U.S. rate of 8.7.

## Syphilis: P&S Rates and Congenital Cases, Idaho, 2004–2013



## Total P&S Syphilis Rates by County, Idaho, 2013



Rate (per 100,000 population)

≤0.4

0.41 - 2.2

>2.2